

LA-UR-12-22825

Approved for public release; distribution is unlimited.

Title:	Preequilibrium Emission of Light Fragments in Spallation Reactions
Author(s):	Kerby, Leslie M. Mashnik, Stepan G. Sierk, Arnold John
Intended for:	12th Annual Student Symposium, LANL, 2012-08-07/2012-08-08 (Los Alamos, New Mexico, United States)



Disclaimer:

Los Alamos National Laboratory, an affirmative action/equal opportunity employer, is operated by the Los Alamos National Security, LLC for the National Nuclear Security Administration of the U.S. Department of Energy under contract DE-AC52-06NA25396. By approving this article, the publisher recognizes that the U.S. Government retains nonexclusive, royalty-free license to publish or reproduce the published form of this contribution, or to allow others to do so, for U.S. Government purposes. Los Alamos National Laboratory requests that the publisher identify this article as work performed under the auspices of the U.S. Department of Energy. Los Alamos National Laboratory strongly supports academic freedom and a researcher's right to publish; as an institution, however, the Laboratory does not endorse the viewpoint of a publication or guarantee its technical correctness.

Precompound Emission of Light Fragments in Spallation Reactions

Leslie M. Kerby, Stepan G. Mashnik, and Arnold J. Sierk

Summer 2012
Los Alamos National Laboratory

Why This Research Is Important



Single Event Upsets (SEUs)

•October 2008, Airbus en route from Perth to Singapore¹

¹Necia Grant Cooper, "The Invisible Neutron Threat", National Security Science Feb. 2012: 13.

•Cold war satellite malfunctioned, detected nuclear missile launch²

²Countdown to Zero, dir. Lucy Walker, perf. Graham Allison, James Baker III, DVD, Magnolia, 2010.

Why This Research Is Important, cont.

Also Important in

- Radiation Shielding
- Medical applications (proton therapy for cancer)
- Understanding better the mechanisms of nuclear reactions


The 2008-2010 IAEA Benchmark of Spallation Models

- Recommended considering preequilibrium emission (and maybe also coalescence production) of fragments heavier than ^4He ^{3,4}

³S. G. Mashnik et al., “Second Advanced Workshop on Model Codes for Spallation Reactions”, CEA-Saclay, France, 8-11 Feb 2010, LA-UR-10-00510.

⁴S. Leray et al., “Results from the IAEA Benchmark of Spallation Models”, Journal of the Korean Physical Society Vol. 59, No. 2 (2011), 791-796.

Current Capabilities of CEM03.03




Evaporation

Precompound


Machner et al., Phys. Rev. C 73 (2006) 044606.

Overview of CEM Model

A general scheme of CEM/LAQGSM calculation



Flowchart for Preequilibrium Code



Modified Exciton Model of Preequilibrium Emission

The Modified Exciton Model (MEM) used by CEM⁵ calculates Γ_j , the emission width (or probability of emitting particle fragment j) as

$$\Gamma_j(p, h, E) = \int_{V_j^c}^{E-B_j} \lambda_c^j(p, h, E, T) dT \quad (1)$$

where the partial transmission probabilities, λ_c^j , are equal to


$$\lambda_c^j(p, h, E, T) = \frac{2s_j + 1}{\pi^2 \hbar^3} \mu_j \Re(p, h) \frac{\omega(p-1, h, E-B_j-T)}{\omega(p, h, E)} T \sigma_{inv}(T) \quad (2)$$

For complex particles, an extra factor γ_j is introduced:


$$\gamma_j \approx p_j^3 \left(\frac{p_j}{A} \right)^{p_j-1} \quad (3)$$

⁵K. K. Gudima, S. G. Mashnik, and V. D. Toneev, "Cascade-Exciton Model of Nuclear Reactions," Nuclear Phys. A401 (1983) 329-361.

Comparison with Hagiwara, et al.⁶





Preliminary Results



Machner et al., Phys. Rev. C 73 (2006) 044606.

Angle-integrated Cross Section

High-energy tails of light fragments obtained without destroying established cross sections.




UNCLASSIFIED

LA-UR-12-22825

200 MeV p + ^{27}Al (after INC)

**Residual
Nuclei
After INC**










Probability



200 MeV p + ^{197}Au (after INC)

**Residual
Nuclei
After INC**


Probability




2500 MeV p + ^{197}Au (after INC)

**Residual
Nuclei
After INC**

Probability



More Preliminary Results



Bubak et al., *Physical Review, Part C, Nuclear Physics* Vol.76 (2007) 014618.

Summary

CEM Extension

- Extended CEM to include emission of light fragments heavier than ^4He in the preequilibrium stage
- Added a subroutine to print the spectra of light fragments, according to isotope, mass number, or Z number
- Built a module to calculate residual nuclei physical properties, which can be inserted anywhere in the reaction process we want

Bugs

- Several bugs were encountered and fixed

Results

- Preliminary results show much greater ability to describe high-energy tails of LF

Future Work

- Parameterize γ_β , investigate coalescence and Fermi break-up. Upgrade evaporation model

Thank you for your attention!

PISA—Tally and Print Spectra Routine

